

# COMPACT $T(1)$ THEOREM À LA STEIN

ÁRPÁD BÉNYI, GUOPENG LI, TADAHIRO OH, AND RODOLFO H. TORRES

ABSTRACT. We prove a compact  $T(1)$  theorem, involving quantitative estimates, analogous to the quantitative classical  $T(1)$  theorem due to Stein. We also discuss the  $C_c^\infty$ -to- $CMO$  mapping properties of non-compact Calderón-Zygmund operators.

## CONTENTS

1. Introduction	1
2. Calderón-Zygmund operators, $BMO$ , and $CMO$	3
3. $T(1)$ theorem and main result	5
3.1. Classical and compact $T(1)$ theorems	5
3.2. $T(1)$ theorems à la Stein	6
4. Proof of Theorem 1	7
5. Mapping properties of Calderón-Zygmund operators into $CMO$	12
Appendix A. Lack of compactness of convolution operators in $L^p(\mathbb{R}^d)$	16
Appendix B. $BMO$ , $VMO$ , and $CMO$ are not weakly sequentially complete	17
References	20

## 1. INTRODUCTION

The classical  $T(1)$  theorem of David and Journé [10] is a fundamental result in harmonic analysis that characterizes the  $L^2$ -boundedness of operators with Calderón-Zygmund kernels. A remarkable insight of this theorem is that one only needs to test whether the actions (properly defined) of the operator and its formal transpose on the constant function 1 belong to  $BMO$  and to verify the so-called weak boundedness property.

There are other equally useful versions of the  $T(1)$  theorem that avoid the introduction of the weak boundedness property. For example, in [10], the  $L^2$ -boundedness is also shown to be equivalent to the uniform boundedness (with respect to  $\xi$ ) in  $BMO$  of the operator and its transpose acting on the character functions  $x \mapsto e^{ix \cdot \xi}$ . In a very elegant formulation, Stein [28] proved a quantitative version of the  $T(1)$  theorem that entirely avoids à priori mentioning  $BMO$  in the statement of the theorem or defining the operator on smooth  $L^\infty$ -functions. He instead considered appropriate uniform  $L^2$ -estimates for the operator and its transpose acting on normalized bump functions; see Theorem C below. Although the actual proof of Stein's version does use the definition of  $T(1)$  and equivalent formulations of the weak boundedness

---

2020 *Mathematics Subject Classification.* 42B20, 47B07.

*Key words and phrases.*  $T(1)$  theorem; Calderón-Zygmund operator; compactness, bounded mean oscillation, vanishing mean oscillation, continuous mean oscillation,  $BMO$ ,  $VMO$ ,  $CMO$ .

property, the advantage of his formulation is that, once the theorem has been established, the definitions of such concepts do not need to be considered when applying the theorem to specific applications. See also [4] for a further discussion on this result by Stein.

Much more recently, several authors have considered versions of the  $T(1)$  theorem characterizing the situation when operators with Calderón-Zygmund kernels are not only bounded but also compact. Such operators do exist and, beyond some simple ones defined on compact sets whose kernels have integrable singularities, an interesting example is provided by a certain class of pseudodifferential operators in  $\mathbb{R}^d$ ; see Cordes [9]. See also [7] for a version of the result for pseudodifferential operators even in a weighted setting. Other examples are provided by appropriate paraproducts, but such operators are usually employed in the proof of compact versions of the  $T(1)$  theorem and thus their compactness need to be proved by different methods.

In [30], Villarroya proved a compactness version of the  $T(1)$  theorem by replacing the weak boundedness property and the condition  $T(1), T^*(1) \in BMO$  in the classical  $T(1)$  theorem (see Theorem A) with an appropriate weak compactness property and the stronger requirement that  $T(1)$  and  $T^*(1)$  belong to  $CMO \subsetneq BMO$  ( $CMO$  denotes the space of functions of *continuous mean oscillation*; see Definition 2.2 below), and by imposing some additional conditions on the kernel of the operator. More recently, Mitkovski and Stockdale [23] in the linear case and Fragkos, Green, and Wick [13] in the multilinear setting, established compactness versions of the  $T(1)$  theorem, avoiding the additional conditions on the kernel of a Calderón-Zygmund operator that appeared in the original version of Villarroya. We note that the space  $CMO$  is a celebrity in harmonic analysis in its own right, appearing naturally in compactness results ever since the classical compactness result of Uchiyama [29] on commutators of Calderón-Zygmund operators with  $CMO$  functions.

The main goal of this article is to present a version of a compact  $T(1)$  theorem in the spirit of Stein's  $T(1)$  theorem on boundedness (Theorem C), without any reference to the actions of  $T$  and  $T^*$  on the constant function 1 in the statement of the result. A secondary goal of this article is to discuss the  $C_c^\infty$ -to- $CMO$  mapping properties of non-compact Calderón-Zygmund operators, both of convolution and non-convolution types. For this purpose, we also revisit some delicate topological properties of  $BMO$ -type spaces; see Appendix B.

This article is organized as follows. In Section 2, we go over some background and definitions needed for the remainder of the paper. After recalling the classical  $T(1)$  theorem and its compact counterpart in Subsection 3.1, we state our main result, a compact  $T(1)$  theorem à la Stein (Theorem 1) in Subsection 3.2. Section 4 is dedicated to the proof of Theorem 1. In Section 5, we then consider the  $C_c^\infty$ -to- $CMO$  mapping properties of non-compact Calderón-Zygmund operators. As a byproduct of our analysis, we also present a direct proof that, under the cancellation assumption  $T(1) = T^*(1) = 0$ , a compact Calderón-Zygmund operator is compact on the Hardy space  $H^1$ ; see Proposition 5.3. In the appendices, we discuss non-compactness of convolution operators as well as the fact about  $BMO$  and some of its subspaces not being weakly sequentially complete, which are of interest on their own. The results in the appendices are probably known to a more specialized functional analysis audience or may follow from more abstract facts. Hence, we do not want to claim authorship on them but they are included here for completeness. We provide references when we could locate them in the

literature, but when we could not, we present some essentially self-contained arguments for some results for the benefit of readers.

## 2. CALDERÓN-ZYGMUND OPERATORS, $BMO$ , AND $CMO$

A linear singular integral operator  $T$  is a map, a priori defined and continuous from  $\mathcal{S}(\mathbb{R}^d)$  into  $\mathcal{S}'(\mathbb{R}^d)$  (the usual Schwartz space and its dual), that takes the form

$$T(f)(x) = \int_{\mathbb{R}^d} K(x, y) f(y) dy \quad (2.1)$$

when the point  $x$  is not in the support of  $f$ . Here, we assume that, away from the diagonal  $\Delta = \{(x, y) \in \mathbb{R}^{2d} : x = y\}$ , the distributional kernel  $K$  of  $T$  coincides with a function that is locally integrable on  $\mathbb{R}^{2d} \setminus \Delta$ , which we still call  $K$ . The formal transpose  $T^*$  of  $T$  is defined similarly with the kernel  $K^*$  given by  $K^*(x, y) = K(y, x)$ .

**Definition 2.1.** A locally integrable function  $K$  on  $\mathbb{R}^{2d} \setminus \Delta$  is called a *Calderón-Zygmund kernel* if it satisfies the following two conditions:

(i) For all  $x, y \in \mathbb{R}^d$ , we have

$$|K(x, y)| \lesssim |x - y|^{-d}. \quad (2.2)$$

(ii) There exists  $\delta \in (0, 1]$  such that

$$|K(x, y) - K(x', y)| + |K(y, x) - K(y, x')| \lesssim \frac{|x - x'|^\delta}{|x - y|^{d+\delta}} \quad (2.3)$$

for all  $x, x', y \in \mathbb{R}^d$  satisfying  $|x - x'| < \frac{1}{2}|x - y|$ .

We say that a linear operator  $T$  of the form (2.1) with a Calderón-Zygmund kernel  $K$  is a *Calderón-Zygmund operator* if  $T$  extends to a bounded operator on  $L^{p_0}(\mathbb{R}^d)$  for some  $1 < p_0 < \infty$ . It is well-known [27] that if  $T$  is a Calderón-Zygmund operator, then it is bounded on  $L^p(\mathbb{R}^d)$  for all  $1 < p < \infty$ . Hence, in the following, we restrict our attention to the  $L^2$ -boundedness of such linear operators. It is also well known that a Calderón-Zygmund operator is bounded from  $L^\infty(\mathbb{R}^d)$  to  $BMO(\mathbb{R}^d)$  and also from  $H^1(\mathbb{R}^d)$  to  $L^1(\mathbb{R}^d)$ . Here,  $H^1(\mathbb{R}^d)$  denotes the Hardy space, which is the predual of  $BMO$  [11], and  $BMO$  denotes the space of functions of *bounded mean oscillation*, which we now recall.

**Definition 2.2.** Given a locally integrable function  $f$  on  $\mathbb{R}^d$ , define the  $BMO$ -seminorm by

$$\|f\|_{BMO} = \sup_Q M_Q(f) = \sup_Q \frac{1}{|Q|} \int_Q |f(x) - \text{ave } f| dx, \quad (2.4)$$

where the supremum is taken over all cubes  $Q$  in  $\mathbb{R}^d$  with sides parallel to the coordinate axes,<sup>1</sup> and

$$\text{ave}_Q f = \frac{1}{|Q|} \int_Q f(x) dx. \quad (2.5)$$

Then, we say that  $f$  is of bounded mean oscillation if  $\|f\|_{BMO} < \infty$  and we define  $BMO(\mathbb{R}^d)$  by

$$BMO(\mathbb{R}^d) = \{f \in L^1_{\text{loc}}(\mathbb{R}^d) : \|f\|_{BMO} < \infty\}.$$

---

<sup>1</sup>In the following, it is understood that all the cubes have sides parallel to the coordinate axes.

As usual, the elements of this space need to be considered as equivalent classes of functions modulo additive constants.

We denote by  $C_c^\infty(\mathbb{R}^d)$  (and by  $C_0(\mathbb{R}^d)$ , respectively) the space of  $C^\infty$ -functions with compact supports (continuous functions vanishing at infinity, respectively). Also, we denote by  $(C_c^\infty)_0(\mathbb{R}^d)$  the subspace of  $C_c^\infty(\mathbb{R}^d)$  consisting of mean-zero functions. We will often suppress the underlying space  $\mathbb{R}^d$  from our notation.

The closure of  $C_c^\infty$  in the  $BMO$  topology is called the space of functions of *continuous mean oscillation*, and it is denoted by  $CMO$ . By definition  $CMO$  is a closed subspace of  $BMO$ , where the topology of  $CMO$  is inherited from that of  $BMO$ , induced by the  $BMO$  semi-norm in (2.4).

We also recall the following characterization of the Hardy space (see, for example, [27, Corollary 1 on p. 221]). Let  $d \geq 2$ .<sup>2</sup> Given  $j = 1, \dots, d$ , let  $R_j$  be the Riesz transform defined by  $\widehat{R_j(f)}(\xi) = i \frac{\xi_j}{|\xi|} \widehat{f}(\xi)$ , where  $\widehat{\cdot}$  denotes the Fourier transform. Then, the Hardy space  $H^1$  consists of functions  $f \in L^1$  such that  $R_j f \in L^1$ ,  $j = 1, \dots, d$ , endowed with the norm:

$$\|f\|_{H^1} = \|f\|_{L^1} + \sum_{j=1}^d \|R_j f\|_{L^1}. \quad (2.6)$$

Next, we recall a characterization of  $CMO$ ; see [29, Lemma on p. 166]. We follow here Bourdaud's formulation in [5, Théorème 7 (iii) on p. 1198]. Given  $f \in L^1_{loc}$ , we define

$$\Gamma_1(f) = \lim_{r \rightarrow 0} \left( \sup_{|Q| \leq r} \frac{1}{|Q|} \int_Q |f(x) - \text{ave}_Q f| dx \right), \quad (2.7)$$

where the supremum is taken over all cubes of volume  $|Q| \leq r$  and  $\text{ave}_Q f$  is as in (2.5).

Similarly, we define

$$\Gamma_2(f) = \lim_{r \rightarrow \infty} \left( \sup_{|Q| \geq r} \frac{1}{|Q|} \int_Q |f(x) - \text{ave}_Q f| dx \right), \quad (2.8)$$

where the supremum is taken over all cubes of volume  $|Q| \geq r$ . Lastly, given a cube  $Q \subset \mathbb{R}^d$ , we define

$$\Gamma_3^Q(f) = \limsup_{\substack{x_0 \in \mathbb{R}^d \\ |x_0| \rightarrow \infty}} \left( \frac{1}{|Q|} \int_{Q+x_0} |f(x) - \text{ave}_{Q+x_0} f| dx \right). \quad (2.9)$$

Then, we have the following characterization of  $CMO$  by the quantities  $\Gamma_1$ ,  $\Gamma_2$ , and  $\Gamma_3^Q$ .

**Lemma 2.3.** *Let  $f \in BMO$ . Then,  $f \in CMO$  if and only if*

$$\Gamma_1(f) = \Gamma_2(f) = \Gamma_3^Q(f) = 0 \quad (2.10)$$

for any cube  $Q \subset \mathbb{R}^d$ .

In [5], Lemma 2.3 was stated in terms of the analogues of the quantities  $\Gamma_1$ ,  $\Gamma_2$ , and  $\Gamma_3^Q$ , where cubes are replaced by balls. A straightforward modification of the argument in [5] yields Lemma 2.3, where we work with cubes.

<sup>2</sup>When  $d = 1$ , we replace the Riesz transforms by the Hilbert transform.

### 3. $T(1)$ THEOREM AND MAIN RESULT

**3.1. Classical and compact  $T(1)$  theorems.** In this subsection, we provide a brief discussion on the classical  $T(1)$  theorem (Theorem A) and its compact counterpart (Theorem B). In order to do so, we need a few more definitions.

**Definition 3.1.** We say that a function  $\phi \in C_c^\infty$  is a *normalized bump function* of order  $M$  if  $\text{supp } \phi \subset B_0(1)$  and  $\|\partial^\alpha \phi\|_{L^\infty} \leq 1$  for all multi-indices  $\alpha$  with  $|\alpha| \leq M$ . Here, given  $r > 0$  and  $x \in \mathbb{R}^d$ ,  $B_x(r)$  denotes the ball of radius  $r$  centered at  $x$ .

Given  $x_0 \in \mathbb{R}^d$  and  $R > 0$ , we set

$$\phi^{x_0, R}(x) = \phi\left(\frac{x - x_0}{R}\right). \quad (3.1)$$

The statement of the  $T(1)$  theorem of David and Journé [10] is the following.

**Theorem A** ( $T(1)$  theorem). *Let  $T : \mathcal{S} \rightarrow \mathcal{S}'$  be a linear singular integral operator with a Calderón-Zygmund kernel. Then,  $T$  can be extended to a bounded operator on  $L^2$  if and only if it satisfies the following two conditions:*

(A.i)  *$T$  satisfies the weak boundedness property; there exists  $M \in \mathbb{N} \cup \{0\}$  such that we have*

$$|\langle T(\phi_1^{x_1, R}), \phi_2^{x_2, R} \rangle| \lesssim R^d \quad (3.2)$$

*for any normalized bump functions  $\phi_1$  and  $\phi_2$  of order  $M$ ,  $x_1, x_2 \in \mathbb{R}^d$ , and  $R > 0$ .*

(A.ii)  *$T(1)$  and  $T^*(1)$  are in  $BMO$ .*

Since  $T$  is a priori defined only in  $\mathcal{S}$ , the expressions  $T(1)$  and  $T^*(1)$  need to be interpreted carefully; see (4.2) below.

**Definition 3.2.** We say that a linear singular integral operator  $T : \mathcal{S} \rightarrow \mathcal{S}'$  with a Calderón-Zygmund kernel has the *weak compactness property* if there exists  $M \in \mathbb{N} \cup \{0\}$  such that we have

$$\lim_{|x_0| + R + R^{-1} \rightarrow \infty} R^{-d} |\langle T(\phi_1^{x_0 + x_1, R}), \phi_2^{x_0 + x_2, R} \rangle| = 0 \quad (3.3)$$

for any normalized bump functions  $\phi_1$  and  $\phi_2$  of order  $M$  and  $x_1, x_2 \in \mathbb{R}^d$ .

The following version of the compact counterpart of the  $T(1)$  theorem (Theorem A) was recently proved in [23]; see also [30]. For a version of a compact  $T(b)$  theorem, see [31].

**Theorem B** (compact  $T(1)$  theorem). *Let  $T$  be as in Theorem A. Then,  $T$  can be extended to a compact operator on  $L^2$  if and only if it satisfies the following two conditions:*

(B.i)  *$T$  satisfies the weak compactness property.*

(B.ii)  *$T(1)$  and  $T^*(1)$  are in  $CMO$ .*

We say that a linear singular integral operator  $T$  with a Calderón-Zygmund kernel is a *compact Calderón-Zygmund operator* if it is compact on  $L^2$ .

**Remark 3.3.** If  $T$  is a compact Calderón-Zygmund operator, then it is also compact

- (a) on  $L^p$  for any  $1 < p < \infty$ ; this follows from interpolation of compactness [19],
- (b) from  $L^\infty$  to  $CMO$ ; see [25],

- (c) from the Hardy space  $H^1$  to  $L^1$ ; this follows from (b) applied to  $T^*$  (which is compact on  $L^2$  in view of Lemma 3.5 (ii)) and duality;<sup>3</sup> see [24, p. 1285],
- (d) from  $L^1$  to  $L^{1,\infty}$ ; see [24],
- (e) from  $BMO$  to  $CMO$  ([25]) under the cancellation assumption that  $T(1) = T^*(1) = 0$ ,
- (f) from  $H^1$  to  $H^1$  under the cancellation assumption that  $T(1) = T^*(1) = 0$ ; this claim follows from (e) and duality; see also Proposition 5.3.

**Remark 3.4.** Our definitions of the weak boundedness property (3.2) and the weak compactness property (Definition 3.3) are slightly different from those in [10, 23], but one can easily check that they are equivalent.

We also note that, as for the weak boundedness property, it suffices to verify (3.2) for  $x_1 = x_2$ ; see [18]. However, it does not seem to be the case for the weak compactness property (3.3).

**3.2.  $T(1)$  theorems à la Stein.** We first recall a quantitative formulation of the classical  $T(1)$  theorem due to Stein; see [28, Theorem 3 on p. 294].

**Theorem C** ( $T(1)$  theorem à la Stein). *Let  $T$  be as in Theorem A. Then,  $T$  can be extended to a bounded operator on  $L^2$  if and only if there exists  $M \in \mathbb{N} \cup \{0\}$  such that we have*

$$\|T(\phi^{x_0,R})\|_{L^2} + \|T^*(\phi^{x_0,R})\|_{L^2} \lesssim R^{\frac{d}{2}} \quad (3.4)$$

for any normalized bump function  $\phi$  of order  $M$ ,  $x_0 \in \mathbb{R}^d$ , and  $R > 0$ .

Here, the conditions (A.i) and (A.ii) in Theorem A are replaced by the quantitative estimate (3.4) involving normalized bump functions. In the following, we seek for a suitable replacement of the condition (3.4) for a compact  $T(1)$  theorem à la Stein. Before proceeding further, we first recall the following basic properties of compact operators.

**Lemma 3.5.** *Let  $X$  and  $Y$  be Banach spaces and  $T : X \rightarrow Y$  be a continuous linear operator.*

- (i) *If  $T$  is compact, then  $T$  maps weakly convergent sequences to strongly convergent sequences.*
- (ii) *The operator  $T$  is compact if and only if its transpose  $T^*$  is compact.*

Lemma 3.5 is well known and its proof can be found in many textbook covering the subject; see, for example, Reed and Simons [26, Theorems VI.11 and VI.12 (c)].

We now derive the  $L^2$ - and  $L^\infty$ -conditions for our formulation of a compact  $T(1)$  theorem à la Stein.

**$L^2$ -condition:** We first rewrite the condition (3.4) as

$$\|T(R^{-\frac{d}{2}}\phi^{x_0,R})\|_{L^2} + \|T^*(R^{-\frac{d}{2}}\phi^{x_0,R})\|_{L^2} \lesssim 1.$$

Note that  $R^{-\frac{d}{2}}\phi^{x_0,R}$  converges weakly to 0 in  $L^2$  as  $|x_0| + R + R^{-1} \rightarrow \infty$ . Suppose now that  $T$  is compact on  $L^2$ . Then, from Lemma 3.5 (i), we see that  $R^{-\frac{d}{2}}T(\phi^{x_0,R})$  and  $R^{-\frac{d}{2}}T^*(\phi^{x_0,R})$

---

<sup>3</sup>Although  $L^1$  is not the dual of  $L^\infty$ , an argument using duality can still be applied because the  $L^1$ -norm can be computed by pairing with  $L^\infty$ -functions.

converge strongly to 0 in  $L^2$  as  $|x_0| + R + R^{-1} \rightarrow \infty$ , namely,

$$\lim_{|x_0|+R+R^{-1} \rightarrow \infty} \left( R^{-\frac{d}{2}} \|T(\phi^{x_0,R})\|_{L^2} + R^{-\frac{d}{2}} \|T^*(\phi^{x_0,R})\|_{L^2} \right) = 0. \quad (3.5)$$

**$L^\infty$ -condition:** Let  $T$  be a Calderón-Zygmund operator that is compact on  $L^2$ . From Remark 3.3(b), we see that  $T$  is compact from  $L^\infty$  to  $CMO$ . Let  $\{f_n\}_{n \in \mathbb{N}}$  be a weak-\* convergent sequence in  $L^\infty$ .<sup>4</sup> Then, it follows from the Banach-Steinhaus theorem that the sequence  $\{f_n\}_{n \in \mathbb{N}}$  is bounded in  $L^\infty$ , and, hence, from the compactness of  $T$  from  $L^\infty$  to  $CMO$ , we see that  $\{T(f_n)\}_{n \in \mathbb{N}}$  is precompact in  $CMO$ . In view of Lemma 3.5(ii), the same conclusion holds for  $T^*$ . This leads to the  $L^\infty$ -condition (ii) in Theorem 1 below.

We are now ready to state our main result, a compact counterpart of the  $T(1)$  theorem à la Stein.

**Theorem 1** (compact  $T(1)$  theorem à la Stein). *Let  $T$  be as in Theorem A. Then,  $T$  can be extended to a compact operator on  $L^2$  if and only if the following two conditions hold:*

- (i) ( $L^2$ -condition). *There exists  $M \in \mathbb{N} \cup \{0\}$  such that (3.5) holds for any normalized bump function  $\phi$  of order  $M$ .*
- (ii) ( $L^\infty$ -condition). *Given any weak-\* convergent sequence  $\{f_n\}_{n \in \mathbb{N}}$  in  $L^\infty$  such that  $\{T(f_n)\}_{n \in \mathbb{N}}$  and  $\{T^*(f_n)\}_{n \in \mathbb{N}}$  are bounded in  $CMO$ , both the sequences  $\{T(f_n)\}_{n \in \mathbb{N}}$  and  $\{T^*(f_n)\}_{n \in \mathbb{N}}$  are precompact in  $CMO$ .*

Note that in the derivation of the  $L^\infty$ -condition and the  $L^2$ -condition before the statement of Theorem 1, we have already established their necessity, assuming that  $T$  is compact. Hence, it remains to prove their sufficiency. Our proof of Theorem 1 follows the lines in the proof of [4, Theorem 1] on the classical  $T(1)$  theorem à la Stein for boundedness of linear singular integral operators. The main task is to verify the conditions (B.i) and (B.ii) in Theorem B by assuming (i) and (ii) in Theorem 1. In particular, for verifying the condition (B.ii), we make use of the characterization of  $CMO$  functions given in Lemma 2.3 and the condition (ii) in Theorem 1.

**Remark 3.6.** It is not clear at this point if the  $L^\infty$ -condition (ii) in Theorem 1 can be removed. See Remark 4.1 below.

**Remark 3.7.** In view of a compact bilinear  $T(1)$  theorem in [13] and other properties of compact bilinear operators recently studied in [3], it would be of interest to investigate an analogue of Theorem 1 in the bilinear setting. We leave this to interested readers.

#### 4. PROOF OF THEOREM 1

In this section, we present a proof of Theorem 1. As mentioned above, it suffices to prove sufficiency of the conditions (i) and (ii). In the following, by assuming the conditions (i) and (ii) in Theorem 1, we verify the conditions (B.i) and (B.ii) in Theorem B, which will in turn imply that  $T$  is compact on  $L^2$ .

---

<sup>4</sup>Recall that  $L^\infty(\mathbb{R}^d) = (L^1(\mathbb{R}^d))^*$ .

We first prove (B.i): the weak compactness property. Let  $\phi_1$  and  $\phi_2$  be normalized bump functions of order 0. From (3.1), we have

$$\|R^{-\frac{d}{2}}\phi_1^{x_0+x_1,R}\|_{L^2} + \|R^{-\frac{d}{2}}\phi_2^{x_0+x_2,R}\|_{L^2} \lesssim 1, \quad (4.1)$$

uniformly in  $x_0, x_1, x_2 \in \mathbb{R}^d$  and  $R > 0$ . Then, from (3.5) and (4.1), we have

$$\begin{aligned} & \lim_{|x_0|+R+R^{-1} \rightarrow \infty} R^{-d} |\langle T(\phi_1^{x_0+x_1,R}), \phi_2^{x_0+x_2,R} \rangle| \\ & \leq \lim_{|x_0|+R+R^{-1} \rightarrow \infty} \min \left( \|T(R^{-\frac{d}{2}}\phi_1^{x_0+x_1,R})\|_{L^2} \|R^{-\frac{d}{2}}\phi_2^{x_0+x_2,R}\|_{L^2}, \right. \\ & \quad \left. \|R^{-\frac{d}{2}}\phi_1^{x_0+x_1,R}\|_{L^2} \|T^*(R^{-\frac{d}{2}}\phi_2^{x_0+x_2,R})\|_{L^2} \right) \\ & = 0 \end{aligned}$$

for any  $x_1, x_2 \in \mathbb{R}^d$ . This proves the condition (B.i) in Theorem B.

Next, we prove the condition (B.ii) in Theorem B. In the following, we only show  $T(1) \in CMO$ , since  $T^*(1) \in CMO$  follows analogously by symmetry. Since (3.5) implies the bound (3.4), it follows from Theorem C that  $T(1) \in BMO$ , where  $T(1)$  is interpreted as

$$\langle T(1), g \rangle = \lim_{R \rightarrow \infty} \langle T(\phi_R), g \rangle \quad (4.2)$$

for any  $g \in (C_c^\infty)_0$ .<sup>5</sup> Here,  $\phi$  is a test function in  $C_c^\infty$  with  $0 \leq \phi \leq 1$  such that  $\phi(x) = 1$  for  $|x| \leq \frac{1}{2}$  and  $\text{supp } \phi \subset B_0(1)$ , and

$$\phi_R(x) = \phi(R^{-1}x). \quad (4.3)$$

Moreover, proceeding as in [4, Section 4] with (3.4), we have

$$\|T(\phi_R)\|_{BMO} \lesssim 1, \quad (4.4)$$

uniformly in  $R > 0$ ; see [4, (4.6)] with  $b_1 = 1$ .

In the following, we show that  $T(\phi_R)$  indeed belongs to  $CMO$  for each  $R \gg 1$  by verifying (2.10) in Lemma 2.3. Fix  $M \in \mathbb{N}$ . Then, by imposing that

$$\|\partial^\alpha \phi\|_{L^\infty} \leq 1 \quad (4.5)$$

for all multi-indices  $\alpha$  with  $|\alpha| \leq M$ , the function  $\phi$  defined above is a normalized bump function of order  $M$ .

Fix a cube  $Q = Q(\ell, x_0)$  of side length  $\ell > 0$  with center  $x_0 \in \mathbb{R}^d$ . Set

$$r = 6 \text{diam}(Q) = 6\sqrt{d}\ell. \quad (4.6)$$

• **Case 1:**  $r \rightarrow \infty$ .

By Cauchy-Schwarz's inequality with (3.5) and (4.6), we have

$$\text{ave}_Q |T(\phi_R)| = \frac{1}{|Q|} \int_Q |T(\phi_R)(x)| dx \leq \frac{1}{|Q|^{\frac{1}{2}}} \|T(\phi_R)\|_{L^2} \lesssim \frac{R^{\frac{d}{2}}}{r^{\frac{d}{2}}} \rightarrow 0,$$

as  $r \rightarrow \infty$ , from which we obtain

$$\frac{1}{|Q|} \int_Q |T(\phi_R) - \text{ave}_Q T(\phi_R)| dx \leq 2 \text{ave}_Q |T(\phi_R)| \rightarrow 0,$$

<sup>5</sup>Recall that  $(C_c^\infty)_0$  is dense in the Hardy space  $H^1$ , which is a predual of  $BMO$ ; see [10, p. 372]. Namely,  $T(1)$  is defined as the weak-\* limit (in  $BMO$ ) of  $T(\phi_R)$ .

as  $r \rightarrow \infty$ . In view of (2.8), this shows

$$\Gamma_2(T(\phi_R)) = \lim_{r \rightarrow \infty} \left( \sup_{|Q| \geq r} \frac{1}{|Q|} \int_Q |T(\phi_R)(x) - \text{ave}_Q T(\phi_R)| dx \right) = 0. \quad (4.7)$$

• **Case 2:**  $|x_0| \rightarrow \infty$ .

Fix  $R, r > 0$ . With  $\phi_Q = \phi^{x_0, r}$  as in (3.1), write  $T(\phi_R)$  as

$$T(\phi_R) = T(\phi_Q \phi_R) + T((1 - \phi_Q)\phi_R) =: \text{I} + \text{II}. \quad (4.8)$$

By choosing  $|x_0| \gg R$ , we have  $\phi_Q \phi_R = 0$  and thus  $\text{I} = 0$ . Hence, we only need to estimate the second term  $\text{II}$  on the right-hand side of (4.8). From the support condition

$$\text{supp}(1 - \phi_Q) \subset \mathbb{R}^d \setminus B_{x_0}(3 \text{diam}(Q)) \subset \mathbb{R}^d \setminus Q$$

and (2.1), we have

$$\text{II}(x) = \int_{\mathbb{R}^d} K(x, y)(1 - \phi_Q(y))\phi_R(y)dy \quad (4.9)$$

for any  $x \in Q$ . By taking an average over  $Q$ , we then have

$$\text{ave}_Q \text{II} = \frac{1}{|Q|} \int_Q \int_{\mathbb{R}^d} K(z, y)(1 - \phi_Q(y))\phi_R(y)dydz. \quad (4.10)$$

By taking the difference of (4.9) and (4.10), we have

$$\text{II}(x) - \text{ave}_Q \text{II} = \frac{1}{|Q|} \int_Q \int_{\mathbb{R}^d} (K(x, y) - K(z, y))(1 - \phi_Q(y))\phi_R(y)dydz$$

for any  $x \in Q$ . In particular, from (2.3), we have

$$\begin{aligned} & |\text{II}(x) - \text{ave}_Q \text{II}| \\ & \lesssim \frac{1}{|Q|} \int_Q \int_{|x-z| \leq \text{diam}(Q) < \frac{1}{2}|x-y|} \frac{|x-z|^\delta}{|x-y|^{d+\delta}} |1 - \phi_Q(y)| |\phi_R(y)| dydz, \end{aligned} \quad (4.11)$$

uniformly in  $x \in Q$ .

By considering the support of  $\phi_R$ , we have  $|y| \lesssim R$  in (4.11). Thus, for  $|x_0| \gg R + r$ , it follows from  $|x - x_0| \leq r$  that  $|x - y| \sim |x| \sim |x_0|$ . Then, by performing the  $y$ -integration in (4.11) on  $\text{supp}(\phi_R)$ , we obtain

$$|\text{II}(x) - \text{ave}_Q \text{II}| \lesssim \frac{r^\delta R^d}{|x_0|^{d+\delta}} \rightarrow 0,$$

as  $|x_0| \rightarrow \infty$ , uniformly in  $x \in Q$ . Hence, together with  $\text{I} = 0$  for  $|x_0| \gg R$ , we obtain

$$\frac{1}{|Q|} \int_Q |T(\phi_R) - \text{ave}_Q T(\phi_R)| dx \rightarrow 0, \quad (4.12)$$

as  $|x_0| \rightarrow \infty$ .

Let  $\tilde{Q} = \tilde{Q}(\ell, x_1)$  be a cube of side length  $\ell > 0$  with center  $x_1 \in \mathbb{R}^d$ . Then, given  $\tilde{x}_0 \in \mathbb{R}^d$ , we have

$$\tilde{Q} + \tilde{x}_0 = Q(\ell, x_0),$$

where  $Q(\ell, x_0)$  denotes the cube of side length  $\ell > 0$  with center  $x_0 := x_1 + \tilde{x}_0$ . For fixed  $x_1 \in \mathbb{R}^d$ , we have  $|x_0| \rightarrow \infty$  as  $|\tilde{x}_0| \rightarrow \infty$ . Therefore, from (2.9) and (4.12), we obtain

$$\Gamma_3^{\tilde{Q}}(T(\phi_R)) = \limsup_{\substack{\tilde{x}_0 \in \mathbb{R}^d \\ |\tilde{x}_0| \rightarrow \infty}} \left( \frac{1}{|\tilde{Q}|} \int_{\tilde{Q} + \tilde{x}_0} |T(\phi_R)(x) - \underset{\tilde{Q} + \tilde{x}_0}{\text{ave}} T(\phi_R)| dx \right) = 0 \quad (4.13)$$

for any cube  $\tilde{Q} \subset \mathbb{R}^d$ .

• **Case 3:**  $r \rightarrow 0$ .

Fix  $R > 0$  and  $x_0 \in \mathbb{R}^d$ . In view of (3.5), we have

$$\lim_{r \rightarrow 0} r^{-\frac{d}{2}} \|T(\phi^{x_0, r})\|_{L^2} = 0.$$

Then, by setting  $A_r = \min(r^{\frac{1}{2}} \|T(\phi^{x_0, r})\|_{L^2}^{-\frac{1}{d}}, r^{-\frac{1}{d+2}})$ , we have

$$A_r \rightarrow \infty, \quad (4.14)$$

$$A_r^{\frac{d+2}{2}} r \rightarrow 0, \quad (4.15)$$

and

$$A_r^{\frac{d}{2}} r^{-\frac{d}{2}} \|T(\phi^{x_0, r})\|_{L^2} \rightarrow 0 \quad \text{or equivalently} \quad r^{-\frac{d}{2}} \|T(\phi^{x_0, A_r r})\|_{L^2} \rightarrow 0, \quad (4.16)$$

as  $r \rightarrow 0$ , where, in (4.16), we used the fact that  $A_r r \rightarrow 0$  as  $r \rightarrow 0$ . In the following, we assume  $r > 0$  is sufficiently small such that  $A_r > 1$ .

We only consider the case  $x_0 \in \text{supp } \phi_R$ , since the case  $x_0 \notin \text{supp } \phi_R$  can be treated in an analogous but easier manner. Define  $\phi_Q$  by

$$\tilde{\phi}_Q = \phi^{x_0, A_r r}. \quad (4.17)$$

Then, we can write  $T(\phi_R)$  as

$$\begin{aligned} T(\phi_R) &= \phi_R(x_0)T(\tilde{\phi}_Q) + T(\tilde{\phi}_Q(-\phi_R(x_0) + \phi_R)) + T((1 - \tilde{\phi}_Q)\phi_R) \\ &=: \tilde{\text{I}} + \tilde{\text{II}} + \tilde{\text{III}}. \end{aligned} \quad (4.18)$$

From (4.17) and (4.16), we have

$$\underset{Q}{\text{ave}} |\tilde{\text{I}}| \leq \frac{|\phi_R(x_0)|}{|Q|} \int_Q |T(\tilde{\phi}_Q)| dx \lesssim \frac{1}{r^{\frac{d}{2}}} \|T(\phi^{x_0, A_r r})\|_{L^2} \rightarrow 0, \quad (4.19)$$

as  $r \rightarrow 0$ .

By the mean value theorem with (4.3), (4.5), and (4.17) (which implies  $|x - x_0| \lesssim A_r r$  on  $\text{supp } \tilde{\phi}_Q$ ), we have

$$|\tilde{\phi}_Q(x)(-\phi_R(x_0) + \phi_R(x))| \lesssim R^{-1} |\tilde{\phi}_Q(x)| |x - x_0| \lesssim R^{-1} A_r r |\tilde{\phi}_Q(x)|. \quad (4.20)$$

Then, from the boundedness of  $T$  on  $L^2(\mathbb{R}^d)$ , (4.20), (4.17), and (4.15), we have

$$\underset{Q}{\text{ave}} |\tilde{\text{III}}| \leq \frac{1}{|Q|^{\frac{1}{2}}} \|\tilde{\text{III}}\|_{L^2(Q)} \lesssim \frac{A_r r}{R r^{\frac{d}{2}}} \|\tilde{\phi}_Q\|_{L^2(\mathbb{R}^d)} \sim \frac{A_r^{\frac{d+2}{2}} r}{R} \rightarrow 0, \quad (4.21)$$

as  $r \rightarrow 0$ .

As for  $\tilde{\mathbb{M}}$ , proceeding as in Case 2, we have

$$\tilde{\mathbb{M}}(x) - \text{ave}_Q \tilde{\mathbb{M}} = \frac{1}{|Q|} \int_Q \int_{\mathbb{R}^d} (K(x, y) - K(z, y)) (1 - \tilde{\phi}_Q(y)) \phi_R(y) dy dz$$

for any  $x \in Q$ . Then, from (2.3) with  $A_r > 1$ , we have

$$\begin{aligned} & |\tilde{\mathbb{M}}(x) - \text{ave}_Q \tilde{\mathbb{M}}| \\ & \lesssim \frac{1}{|Q|} \int_Q \int_{|x-z| \leq \text{diam}(Q) < \frac{1}{2A_r}|x-y|} \frac{|x-z|^\delta}{|x-y|^{d+\delta}} |1 - \phi_Q(y)| |\phi_R(y)| dy dz, \end{aligned}$$

uniformly in  $x \in Q$ . By integrating in  $y$  and  $z$  in the polar coordinates (centered at  $x$  with  $r_1 = |x - y| \gtrsim A_r r$  and  $r_2 = |x - z| \lesssim r$ ) with  $|Q| \sim r^d$  and (4.14), we have

$$|\tilde{\mathbb{M}}(x) - \text{ave}_Q \tilde{\mathbb{M}}| \lesssim \frac{1}{r^d} \int_0^{c_2 r} \int_{c_1 A_r r}^\infty \frac{1}{r_1^{1+\delta}} dr_1 r_2^{d-1+\delta} dr_2 \sim \frac{1}{A_r^\delta} \longrightarrow 0, \quad (4.22)$$

as  $r \rightarrow 0$ , uniformly in  $x \in Q$ .

Hence, putting (4.18), (4.19), (4.21), and (4.22) together,

$$\frac{1}{|Q|} \int_Q |T(\phi_R) - \text{ave}_Q T(\phi_R)| dx \longrightarrow 0, \quad (4.23)$$

as  $r \rightarrow 0$ , uniformly in  $R \gg 1$ . From the definition (2.7) of  $\Gamma_1$ , this shows

$$\Gamma_1(T(\phi_R)) = \lim_{r \rightarrow 0} \left( \sup_{|Q| \leq r} \frac{1}{|Q|} \int_Q |T(\phi_R)(x) - \text{ave}_Q T(\phi_R)| dx \right) = 0. \quad (4.24)$$

Therefore, from Lemma 2.3 with (4.7), (4.13), and (4.24), we conclude that

$$T(\phi_R) \in CMO \quad (4.25)$$

for each  $R > 0$ .

Let  $\{R_j\}_{j \in \mathbb{N}}$  be an increasing sequence of positive numbers such that  $R_j \rightarrow \infty$  as  $j \rightarrow \infty$ . Then, the sequence  $\{\phi_{R_j}\}_{j \in \mathbb{N}}$  converges to the constant function  $\phi(0) = 1$  in the weak-\* topology of  $L^\infty$ . Moreover, from (4.4) and (4.25), the sequence  $\{T(\phi_{R_j})\}_{j \in \mathbb{N}}$  is bounded in  $CMO$ . Therefore, from the  $L^\infty$ -condition (ii), we conclude that there exists a subsequence  $\{T(\phi_{R_{j_k}})\}_{k \in \mathbb{N}}$  converging strongly to  $T(1)$  in  $CMO$ , which in particular implies  $T(1) \in CMO$ .

We have now verified (B.i) and (B.ii) in Theorem B, which in turn implies that  $T$  is compact on  $L^2$ . This concludes the proof of Theorem 1.

**Remark 4.1.** In view of (4.23), the convergence in (4.24) is uniform in  $R \gg 1$ . However, the convergence in (4.7) and (4.13) in Cases 1 and 2, respectively, is not uniform in  $R \gg 1$ . This is the reason that we needed to introduce the  $L^\infty$ -condition (ii) in Theorem 1.

**Remark 4.2.** In view of  $(CMO)^* = H^1$  (see [8, Theorem (4.1) on p.638]<sup>6</sup>) and that  $T(\phi_R) \in CMO$ ,  $R > 0$ , it follows from (4.2) (which holds for any  $g \in H^1$ ) that  $T(1)$  is a weak limit of  $CMO$  functions with respect to the weak topology on  $CMO$ . Hence, if  $CMO$  were weakly sequentially complete, we would be able to conclude  $T(1) \in CMO$  without the  $L^\infty$ -condition (ii) in Theorem 1. However, this is not the case. See Appendix B.

<sup>6</sup>We note that the space  $VMO$  in [8] corresponds to  $CMO$  in the current work.

5. MAPPING PROPERTIES OF CALDERÓN-ZYGMUND OPERATORS INTO  $CMO$ 

While every Calderón-Zygmund operator maps  $L^\infty$  to  $BMO$ , compact ones in particular map  $L^\infty$  to  $CMO$  as pointed out in Remark 3.3 (b). We will show that, in general, without the compactness assumption, Calderón-Zygmund operators fail to have this property. Moreover, this failure persists even if we restrict input functions to smaller subspaces of  $L^\infty$ , consisting of smooth functions; see Proposition 5.5.

In the case of Calderón-Zygmund operators of convolution type, we do have that they map  $C_0$  into  $CMO$ . This claim appears in [28, p. 180]<sup>7</sup> without a proof. In Proposition 5.4, we prove this claim in a little more generality. We note that, in [16, p. 207] and [23, p. 10], analogous claims are made for general Calderón-Zygmund operators (in particular for those of non-convolution type), which are unfortunately incorrect. In Proposition 5.5, we construct an example of a Calderón-Zygmund operator  $T$  such that  $T(C_c^\infty) \not\subset CMO$ .

We first recall some further properties of Calderón-Zygmund operators. Given a Calderón-Zygmund operator  $T$ , we have  $T(f) \in L^1$  for  $f \in (C_c^\infty)_0$ . Here,  $(C_c^\infty)_0$  denotes the subspace of  $C_c^\infty$  consisting of mean-zero functions, which is dense in  $H^1$ . If, in addition,  $T$  maps  $H^1$  into itself, then we have

$$0 = \int_{\mathbb{R}^d} T(f)(x) dx = \langle 1, T(f) \rangle = \langle T^*(1), f \rangle$$

for any  $f \in (C_c^\infty)_0$ . Namely, we have  $T^*(1) = 0$ . In fact, such a condition is sufficient for boundedness of  $T$  on  $H^1$ .

**Lemma 5.1.** *Let  $T$  be a Calderón-Zygmund operator of convolution type<sup>8</sup> or, more generally, let  $T$  be a Calderón-Zygmund operator with  $T^*(1) = 0$ . Then,  $T$  maps  $H^1$  into itself.*

In the convolution case, this result goes back to Fefferman and Stein [11, Corollary 1]. For non-convolution operators with  $T^*(1) = 0$ , this result was proved by Alvarez and Milman [1, Theorem 1.1]. Under the additional assumption that  $T(1) = 0$ , this result was shown by Frazier et al. [14, 15]. This case subsumes the Fefferman-Stein result on Calderón-Zygmund operators of convolution type; see, for example, the comments in [15, p. 67]. Note that if  $T(1) = T^*(1) = 0$ , then we have both  $T, T^* : H^1 \rightarrow H^1$  and  $BMO \rightarrow BMO$ . The next two results show that the cancellation condition  $T(1) = T^*(1) = 0$  also provides additional mapping properties.

**Lemma 5.2.** *Let  $T$  be a Calderón-Zygmund operator of convolution type or, more generally, let  $T$  be a Calderón-Zygmund operator with  $T(1) = T^*(1) = 0$ . Then, for any  $f \in C_c^\infty$ , we have  $T(f) \in C_0$ .*

*Proof.* We note that this is very easy to prove in the convolution case. Since  $T$  is bounded on  $L^2$ , there exists  $m \in L^\infty$  such that  $\widehat{T(f)} = m\widehat{f}$ . Since  $f \in C_c^\infty$ , we have  $\widehat{f} \in \mathcal{S}$  and thus  $m\widehat{f} \in L^1$ . Then, by the Riemann-Lebesgue lemma, we obtain  $T(f) \in C_0$ .

In the more general case, we can not take such direct advantage of the Fourier transform. Using certain molecular decompositions, Meyer [21, Proposition 2] showed that, under the

<sup>7</sup>The space  $VMO$  in [28] also corresponds to  $CMO$  in our discussion.

<sup>8</sup>Namely, the kernel  $K$  of the operator  $T$  is of the form  $K(x, y) = k(x - y)$  such that  $Tf = k * f$ , at least in the distributional sense. Note that, for a Calderón-Zygmund operator  $T$  of convolution type, we have  $T(1) = T^*(1) = 0$  (as elements in  $BMO$ , namely, modulo additive constants).

hypothesis of the lemma,  $T$  maps Hölder continuous functions into Hölder continuous functions. In particular, for  $f \in C_c^\infty$ ,  $T(f)$  is continuous.

Suppose that  $\text{supp } f \subset B_0(R) \subset \mathbb{R}^d$ . Then, if  $|x| \geq 2R$ , we have  $|x - y| \sim |x|$  for any  $y \in \text{supp } f$ . Hence, from (2.2), we have

$$|T(f)(x)| = \left| \int_{\text{supp } f} K(x, y) f(y) dy \right| \lesssim \int_{\text{supp } f} \frac{1}{|x|^d} |f(y)| dy \lesssim \frac{\|f\|_{L^1}}{|x|^d} \longrightarrow 0 \quad (5.1)$$

as  $|x| \rightarrow \infty$ . Therefore, we conclude that  $T(f) \in C_0$ .  $\square$

As stated in Remark 3.3 (e) and (f), under the cancellation condition  $T(1) = T^*(1) = 0$ , a compact Calderón-Zygmund operator  $T$  is compact from  $BMO$  to  $CMO$  and also from  $H^1$  into itself, where the latter claim follows from the former via duality. In fact, a compact Calderón-Zygmund operator  $T$  is compact from  $CMO$  to itself and thus from Lemma 3.5 (ii), we see that  $T$  is compact from  $H^1$  into itself. Nonetheless, using the characterization (2.6) of the Hardy space  $H^1$ , we present a direct proof of the compactness on  $H^1$  of a compact Calderón-Zygmund operator  $T$  under the cancellation condition  $T(1) = T^*(1) = 0$  without using such a duality argument.

**Proposition 5.3.** *Let  $T$  be a Calderón-Zygmund operator of convolution type or, more generally, let  $T$  be a Calderón-Zygmund operator with  $T(1) = T^*(1) = 0$ . Suppose that  $T$  is compact on  $L^2$ . Then,  $T$  is compact from  $H^1$  to itself.*

*Proof.* Let  $T$  be a compact Calderón-Zygmund operator with  $T(1) = T^*(1) = 0$ . By Remark 3.3 (c), we know that  $T : H^1 \rightarrow L^1$  is compact. On the other hand, recalling from [22, Section 8.9] that the collection of Calderón-Zygmund operators  $S$  satisfying the cancellation condition  $S(1) = S^*(1) = 0$  forms an algebra, we see that  $R_j T$  is also a Calderón-Zygmund operator,  $j = 1, \dots, d$  (since  $R_j(1) = R_j^*(1) = 0$  as a convolution operator). Moreover, since  $T$  is compact on  $L^2$  and  $R_j$  is bounded on  $L^2$ , the composition  $R_j T$  is compact on  $L^2$ . Therefore, the desired compactness on  $H^1$  now follows from the definition (2.6) of the  $H^1$ -norm:

$$\|Tf\|_{H^1} = \|Tf\|_{L^1} + \sum_{j=1}^d \|(R_j T)f\|_{L^1},$$

since each operator appearing on the right-hand side is compact from  $H^1$  into  $L^1$ .  $\square$

We now state the first main result of this section which extends the aforementioned claim in [28, p. 180] to a slightly general setting.

**Proposition 5.4.** *Let  $T$  be a Calderón-Zygmund operator of convolution type or, more generally, let  $T$  be a Calderón-Zygmund operator with  $T(1) = T^*(1) = 0$ . Then,  $T$  maps  $C_0$  to  $CMO$ .*

*Proof.* Let  $f \in C_0$ . Then, there exists  $\{f_n\}_{n \in \mathbb{N}} \subset C_c^\infty$  such that  $f_n$  converges to  $f$  in  $L^\infty$  (and thus in  $BMO$ ) as  $n \rightarrow \infty$ . By the boundedness of  $T : L^\infty \rightarrow BMO$ , we have  $T(f), T(f_n) \in BMO$  for any  $n \in \mathbb{N}$ . Then, from the definition of the transpose and Lemma 5.1, we have

$$\begin{aligned} |\langle T(f) - T(f_n), g \rangle| &= |\langle f - f_n, T^*(g) \rangle| \leq \|f - f_n\|_{BMO} \|T^*(g)\|_{H^1} \\ &\lesssim \|f - f_n\|_{BMO} \|g\|_{H^1} \end{aligned} \quad (5.2)$$

for any  $g \in H^1$ . Recalling

$$\|b\|_{BMO} = \sup_{\|g\|_{H^1}=1} |\langle b, g \rangle|,$$

we see from (5.2) that  $T(f_n)$  converges to  $T(f)$  in  $BMO$  as  $n \rightarrow \infty$ . On the other hand, it follows from Lemma 5.2 that  $T(f_n) \in C_0$  for each  $n \in \mathbb{N}$ . Therefore, recalling from [5, Théorème 7] that the closure of  $C_0$  in  $BMO$  is  $CMO$ , we conclude that  $T(f) \in CMO$ .  $\square$

Our next goal is to show that Proposition 5.4 does not hold in general without the cancellation condition  $T(1) = T^*(1) = 0$ ; see Proposition 5.5. For this purpose, we first need to recall the notion of a paraproduct.

Let  $\varphi, \psi \in C_c^\infty$  be radial functions such that  $\text{supp}(\varphi) \subset B_0(1)$ ,  $\widehat{\psi}(0) = 0$ , and

$$\int_0^\infty |\widehat{\psi}(te_1)|^2 \frac{dt}{t} = 1,$$

where  $e_1 = (1, 0, \dots, 0) \in \mathbb{R}^d$ . For  $t \in \mathbb{R}_+$ , we also define the linear convolution operators  $P_t$  and  $Q_t$  by

$$P_t f = \varphi_t * f \quad \text{and} \quad Q_t f = \psi_t * f, \quad (5.3)$$

where  $h_t = t^{-d}h(t^{-1}\cdot)$  for a function  $h$  on  $\mathbb{R}^d$ . Then, the Calderón reproducing formula [6] states that

$$\int_0^\infty Q_t^2 f \frac{dt}{t} = f$$

holds in  $L^2$ , where  $Q_t^2 f = Q_t(Q_t f) = \psi_t * \psi_t * f$ . Given  $b \in BMO$ , we define a paraproduct  $\Pi_b$  by

$$\Pi_b(f) = \int_0^\infty Q_t((Q_t b)(P_t f)) \frac{dt}{t}, \quad (5.4)$$

a priori defined for  $f \in \mathcal{S}$ . It is well known by now that if  $b \in BMO$  then  $\Pi_b$  is a Calderón-Zygmund operator, satisfying

$$\Pi_b(1) = b \quad \text{and} \quad \Pi_b^*(1) = 0, \quad (5.5)$$

where once again the equalities are understood by testing against functions in  $(C_c^\infty)_0$ ; see, for example, [10, Lemmas 1 and 2]. See also (4.2).

In what follows, we will prove that, for an appropriate symbol  $b \in BMO \setminus CMO$ , the paraproduct  $\Pi_b$  does *not* map  $C_c^\infty$  into  $CMO$ . A naive approach would be to proceed as follows. Fix  $b \in BMO \setminus CMO$ . Assuming that  $\Pi_b$  maps  $C_c^\infty$  into  $CMO$ , we would have  $\Pi_b(\phi_R) \in CMO$ , where  $\phi_R$  is as in (4.3). If  $CMO$  were weakly sequentially complete, we would then have  $b = \Pi_b(1) \in CMO$ , since, as seen in (4.2),  $\Pi_b(1)$  is defined as the limit of  $\Pi_b(\phi_R)$  by testing against  $H^1$ -functions, namely as a limit with respect to the weak topology in  $CMO$ , in view of the duality  $(CMO)^* = H^1$ . This would be a contradiction with the fact that  $b \in BMO \setminus CMO$ . However, this naive approach breaks down precisely because  $CMO$  is *not* sequentially complete; see again Appendix B and Remark 4.2. Therefore, a more delicate argument and a more careful choice of the symbol  $b$  is needed here.

**Proposition 5.5.** *There exists a Calderón-Zygmund operator, in particular a paraproduct  $\Pi_b$  with  $b \in BMO \setminus CMO$  such that  $\Pi_b(C_c^\infty) \not\subset CMO$ .*

We begin by showing that, for our argument, the  $CMO$  space can be “replaced” by  $VMO$ , the space of functions of *vanishing mean oscillation* which is defined as the closure of  $C^\infty \cap BMO$  in the  $BMO$  topology. In particular, we have

$$CMO \subset VMO \subset BMO. \quad (5.6)$$

We recall the following characterization of  $VMO$ ; see, for example, [5, Théorème 5].

**Lemma 5.6.** *Let  $f \in BMO$ . Then,  $f \in VMO$  if and only if  $\Gamma_1(f) = 0$ , where  $\Gamma_1$  is as in (2.7).*

The next lemma shows that  $CMO$  is the subspace of function in  $VMO$  which “vanish” at infinity; see [5, Théorème 7].

**Lemma 5.7.** *Let  $BMO_c$  denote the subspace of  $BMO$  consisting of functions with compact support. Then, we have*

$$CMO = VMO \cap \overline{BMO_c},$$

where the closure is with respect to the  $BMO$ -norm.

Let  $f \in C_c^\infty(\mathbb{R}^d)$ . As we saw in (5.1), we have that  $T(f)$  (which is in  $BMO$ ) vanishes at infinity. Hence, from [5, Théorème 6], we conclude that  $T(f) \in \overline{BMO_c}$ . Therefore, it follows from Lemma 5.7 that  $T(f) \in CMO$  if and only if  $T(f) \in VMO$ .

We are now ready to prove Proposition 5.5.

*Proof of Proposition 5.5.* In the following, we work on  $\mathbb{R}$ . For a suitable choice of  $b \in BMO \setminus VMO$ , we show that  $\Pi_b(\phi_R) \notin VMO$  for any finite  $R \gg 1$  (and hence  $\Pi_b(\phi_R) \notin CMO$  in view of (5.6)), where  $\phi_R$  is as in (4.3).

From (5.3), we have

$$Q_t((Q_t b)(P_t \phi_R))(x) = \frac{1}{t^{2d}} \int_{\mathbb{R}^2} \psi\left(\frac{x-y}{t}\right) Q_t b(y) \varphi\left(\frac{y-z}{t}\right) \phi_R(z) dz dy. \quad (5.7)$$

Suppose that  $\text{supp } \psi, \text{supp } \varphi \subset [-C_0, C_0]$  for some  $C_0 > 0$ . Then, we have  $|x-y|, |y-z| \leq tC_0$  on the right-hand side of (5.7). Thus, by the triangle inequality, we have

$$|z| \leq 2tC_0 + |x| \leq 2C_0 + 1 \quad (5.8)$$

for any  $0 < t \leq 1$  and  $|x| \leq 1$ . Hence, from (5.7) with (4.3) and (5.8), we have

$$Q_t((Q_t b)(P_t \phi_R))(x) = Q_t((Q_t b)(P_t 1))(x) \quad (5.9)$$

for any  $0 < t \leq 1$ ,  $|x| \leq 1$ , and  $R \geq 4C_0 + 2$ .

We now choose a suitable function  $b \in BMO \setminus VMO$ . Define an odd function  $b \in C^\infty((-\infty, 0) \cup (0, \infty); \mathbb{R})$  by setting

$$b(x) = \begin{cases} 1, & \text{for } 0 < x \leq 1, \\ -1, & \text{for } -1 \leq x < 0, \\ 0, & \text{for } x = 0 \text{ and } |x| \geq 2 \end{cases}$$

and smoothly interpolating on the intervals  $[-2, -1]$  and  $[1, 2]$  in such a way that  $b$  is odd. While we have  $b \in L^\infty \subset BMO$ , we see that  $b \notin VMO$  due to the jump at  $x = 0$ . Indeed, with  $Q = [-a, a]$ , we have

$$M_Q(b) = \frac{1}{|Q|} \int_Q |b(x) - \text{ave}_Q b| dx = \frac{1}{|Q|} \int_Q |b(x)| dx = 1$$

for any  $0 < a < 1$ , which implies  $b \notin VMO$  in view of Lemma 5.6.

From (5.4) and (5.5), we have

$$b = \Pi_b(1) = \int_0^\infty Q_t((Q_t b)(P_t 1)) \frac{dt}{t},$$

where the second inequality holds as elements in the dual space of  $H^1$  (namely, as elements in  $BMO$ ). Write  $b = b_1 + b_2$ , where  $b_1$  is defined by

$$b_1 = \int_0^1 Q_t((Q_t b)(P_t 1)) \frac{dt}{t}. \quad (5.10)$$

It is easy to see that  $b_2$  is a smooth function on  $\mathbb{R}$  since it essentially corresponds to the low frequency part of the paraproduct  $\Pi_b(1)$ . Moreover, since  $\varphi$  and  $\psi$  are even and  $b$  is odd, it follows from (5.3) that  $b_2$  is a smooth odd function which in particular implies  $b_2(0) = 0$ . Hence, we conclude that  $b_1$  has a jump discontinuity at  $x = 0$ :

$$b_1(0+) - b_1(0-) = 2. \quad (5.11)$$

Fix  $R \geq 4C_0 + 2$ . Write  $\Pi_b(\phi_R) = A_1 + A_2$ , where  $A_1$  is defined by

$$A_1 = \int_0^1 Q_t((Q_t b)(P_t \phi_R)) \frac{dt}{t}. \quad (5.12)$$

Then, from (5.9), (5.10), and (5.12), we have  $A_1(x) = b_1(x)$  for any  $|x| \leq 1$ . Hence, from (5.11), we obtain

$$A_1(0+) - A_1(0-) = 2,$$

where we used the fact that  $A_2(0) = 0$  just as for  $b_2$  discussed above. By noting that  $A_2$  is a smooth function on  $\mathbb{R}$ , we then have

$$\Pi_b(\phi_R)(0+) - \Pi_b(\phi_R)(0-) = 2.$$

Hence, by computing  $M_Q(\Pi_b(\phi_R))$  for  $Q = (-a, a)$  with  $a$  tending to 0, we see that  $\Gamma_1(\Pi_b(\phi_R)) \neq 0$ , where  $M_Q$  and  $\Gamma_1$  are as in (2.4) and (2.7), respectively. Therefore, we conclude from Lemma 5.6 that  $\Pi_b(\phi_R) \notin VMO$ . This shows that, in general, a Calderón-Zygmund operator does *not* map  $C_c^\infty$  into  $VMO$  or  $CMO$ .  $\square$

**Remark 5.8.** As subspaces of  $BMO$ , the difference between  $CMO$  and  $VMO$  appears only in the global aspect of their elements, namely the decay at spatial infinity of their elements (just like the difference between  $C_0$  and  $C$  as subspaces of  $L^\infty$ ). In the counterexample presented above, the failure of  $C_c^\infty$ -to- $CMO$  mapping property comes from the local property of elements shared by  $CMO$  and  $VMO$ , namely, they can not have an isolated jump discontinuity, while an element in  $BMO$  can have an isolated jump discontinuity,

#### APPENDIX A. LACK OF COMPACTNESS OF CONVOLUTION OPERATORS IN $L^p(\mathbb{R}^d)$

From the result in the previous section, it may be tempting to think that some Calderón-Zygmund operators of convolution type may be compact. However it is known that that cannot be true. See for example, Feitchinger [12, p. 310] where it is stated “that in the case of a non-compact group there do not exist compact multipliers between two isometrically translation invariant spaces”. We will provide a simple proof of this fact in the context of the Lebesgue spaces  $L^p(\mathbb{R}^d)$ .

Consider first the following simple but illustrative example. Given  $\varepsilon > 0$ , let  $J^{-\varepsilon}$  be the Bessel potential of order  $\varepsilon > 0$ . Namely,  $J^{-\varepsilon}$  is a Fourier multiplier operator with a bounded multiplier  $(1 + |\xi|^2)^{-\frac{\varepsilon}{2}}$ . It is not difficult to show that  $J^{-\varepsilon}$  is a Calderón-Zygmund operator. If  $J^{-\varepsilon}$  were compact on  $L^2(\mathbb{R}^d)$ , then this would imply that  $L^2(\mathbb{R}^d)$  compactly embeds into  $H^{-\varepsilon}(\mathbb{R}^d)$ , which is false. However, the situation is different in the compact setting. We point out that if the underlying space is the  $d$ -dimensional torus  $\mathbb{T}^d = (\mathbb{R}/\mathbb{Z})^d$ , then Rellich's lemma states that the embedding  $L^2(\mathbb{T}^d) \hookrightarrow H^{-\varepsilon}(\mathbb{T}^d)$  is indeed compact. Namely, the lack of compactness of the Bessel potential  $J^{-\varepsilon}$  on  $L^2(\mathbb{R}^d)$  comes from the fact that functions on  $\mathbb{R}^d$  can run off to the spatial infinity.

The example above is representative of the following general fact.

**Proposition A.1.** *Let  $1 \leq p < \infty$ . A non-trivial<sup>9</sup> translation invariant operator  $T$  is not a compact operator on  $L^p(\mathbb{R}^d)$ .*

*Proof.* Without loss of generality, we assume that  $T$  is bounded on  $L^p(\mathbb{R}^d)$  since, otherwise,  $T$  is obviously not compact. Since  $T$  is translation invariant, we have

$$T(\tau_y f) = \tau_y T(f), \tag{A.1}$$

where  $\tau_y f(x) = f(x - y)$ .

Fix  $f \in L^p$  such that  $T(f) \neq 0$ . Given a sequence  $\{y_n\}_{n \in \mathbb{N}} \subset \mathbb{R}^d$  with  $|y_n| \rightarrow \infty$ , let  $f_n = \tau_{y_n} f$ . Clearly, the sequence  $\{f_n\}_{n \in \mathbb{N}}$  is bounded in  $L^p$ . If  $T$  were compact on  $L^p$ , then it would follow from the Kolmogorov-Riesz theorem (see [32, Theorem on p. 275]; see also [17, Theorem 5]) that there exists a subsequence  $\{f_{n_k}\}_{k \in \mathbb{N}}$  such that, given any  $\varepsilon > 0$ , there exists  $R = R(\varepsilon) > 0$  such that

$$\int_{|x| > R} |T(f_{n_k})(x)|^p dx < \varepsilon^p \tag{A.2}$$

for any  $k \in \mathbb{N}$ .

Fix  $0 < \varepsilon < \frac{1}{2} \|Tf\|_{L^p}$ . Suppose that there exists  $R = R(\varepsilon) > 0$  such that (A.2) holds. Then, from the definition of  $f_{n_k}$ , (A.1), a change of variables, and the dominated convergence theorem together with the  $L^p$ -boundedness of  $T$ , we have

$$\begin{aligned} \lim_{k \rightarrow \infty} \int_{|x| > R} |T(f_{n_k})(x)|^p dx &= \lim_{k \rightarrow \infty} \int_{|x| > R} |T(f)(x - y_{n_k})|^p dx \\ &= \lim_{k \rightarrow \infty} \int_{|x + y_{n_k}| > R} |T(f)(x)|^p dx = \|Tf\|_{L^p}^p > 2^p \varepsilon^p, \end{aligned}$$

which is a contradiction to (A.2). Therefore, we conclude that a non-trivial translation invariant  $T$  is not compact on  $L^p$ .  $\square$

#### APPENDIX B. $BMO$ , $VMO$ , AND $CMO$ ARE NOT WEAKLY SEQUENTIALLY COMPLETE

In this appendix, we study weakly sequential completeness properties for the  $BMO$ -type spaces which is closely related to the need for the  $L^\infty$ -condition in the proof of Theorem 1; see Remark 4.1 (ii).

Weakly sequential completeness of the Hardy space  $H^1$  follows as an immediate consequence of [8, Lemma (4.2)]. A natural question is then to ask whether the dual and pre-dual of  $H^1$

---

<sup>9</sup>Namely,  $T \neq 0$ .

(*BMO* and *CMO*, respectively) are also weakly sequential complete. We will show that the answer is negative.

**Proposition B.1.** *BMO, VMO, and CMO are not weakly sequentially complete.*

Before proceeding to a proof of Proposition B.1, we state several lemmas. We first recall the following lemma from [2, Proposition 2.3].

**Lemma B.2.** *Let  $X$  be a weakly sequentially complete Banach space. Then, any closed subspace  $Y$  of  $X$  is a weakly sequentially complete Banach space.*

Let  $c_0 = c_0(\mathbb{N}) \subset \ell^\infty(\mathbb{N})$  denote the space of bounded sequences  $\mathbf{a} = \{a_n\}_{n \in \mathbb{N}}$  such that  $\lim_{n \rightarrow \infty} a_n = 0$ . If we endow  $c_0$  with the supremum norm, then we have  $(c_0)^* = \ell^1(\mathbb{N}) = \ell^1$ , where the duality pairing is given by

$$\langle \mathbf{a}, \mathbf{b} \rangle = \sum_{n=1}^{\infty} a_n b_n$$

for  $\mathbf{a} \in c_0$  and  $\mathbf{b} \in \ell^1$ . The space  $c_0$  provides a useful example for many functional analytic situations related to weak sequential (non-)completeness. For the readers convenience, we include a proof of the following fact.

**Lemma B.3.**  *$c_0$  is not weakly sequentially complete.*

*Proof.* Given  $k \in \mathbb{N}$ , define a sequence  $\mathbf{a}_k = \{a_{kn}\}_{n \in \mathbb{N}}$  by setting  $a_{kn} = 1$  for  $n \leq k$  and  $a_{kn} = 0$  for  $n > k$ . Then, it is easy to verify that  $\mathbf{a}_k \in c_0$  for each  $k \in \mathbb{N}$  and that  $\mathbf{a}_k$  is weakly Cauchy. However, its pointwise limit is a constant sequence with value 1 which is not in  $c_0$ . This proves the claim.  $\square$

Given a Banach space  $X$ , we say that  $X$  contains a copy of  $c_0$ , if there exist a sequence  $\{x_n\}_{n \in \mathbb{N}}$  of unit vectors and constants  $C_1, C_2 > 0$  such that

$$C_1 \sup_{n \in \mathbb{N}} |a_n| \leq \left\| \sum_{n=1}^{\infty} a_n x_n \right\|_X \leq C_2 \sup_{n \in \mathbb{N}} |a_n| \quad (\text{B.1})$$

for any  $\mathbf{a} = \{a_n\}_{n \in \mathbb{N}} \in c_0$ .

Let

$$Y = \left\{ x = \sum_{n=1}^{\infty} a_n x_n : \{a_n\}_{n \in \mathbb{N}} \in c_0 \right\}. \quad (\text{B.2})$$

In view of (B.1), we see that every element in  $Y$  is uniquely represented as

$$\mathbf{a} \cdot \mathbf{x} = \sum_{n=1}^{\infty} a_n x_n$$

for some  $\mathbf{a} = \{a_n\}_{n \in \mathbb{N}} \in c_0$ . Moreover,  $Y$  is isomorphic and “almost isometric” to  $c_0$ , meaning that

$$\|\mathbf{a}\|_{c_0} \approx \|\mathbf{a} \cdot \mathbf{x}\|_X$$

in the sense of (B.1). Once again, in view of (B.1), we see that  $Y$  is closed in  $X$  because  $c_0$  is a complete Banach space.

**Lemma B.4.** *Let  $X$  be a Banach space. If  $X$  contains a copy of  $c_0$ , then  $X$  is not weakly sequentially complete.*

Under an additional assumption that the space  $X$  has an unconditional basis, the converse of Lemma B.4 also holds true; see [20, Theorem 1.c.10].

*Proof of Lemma B.4.* Since  $X$  contains a copy of  $c_0$ , there exists  $\{x_n\}_{n \in \mathbb{N}} \subset X$ , satisfying (B.1). Define a subspace  $Y$  of  $X$  as in (B.2). Recalling that  $c_0^* = \ell^1$ , we see that  $Y^* \cong \ell^1$ . Indeed, given  $\mathbf{b} = \{b_n\}_{n \in \mathbb{N}} \in \ell^1$ , we set  $\mathbf{b}(x) = \sum_{n=1}^{\infty} a_n b_n$  for  $x = \mathbf{a} \cdot \mathbf{x} \in Y$ . By noting that the  $Y$ - $Y^*$  duality pairing agrees with the  $c_0$ - $\ell^1$  duality pairing, it follows from Lemma B.3 that  $Y$  is not weakly sequentially complete. Finally, by recalling that  $Y$  is a closed subspace of  $X$ , we conclude from Lemma B.2 that  $X$  is not weakly sequentially complete.  $\square$

We are now ready to present a proof of Proposition B.1.

*Proof of Proposition B.1.* We only consider the  $d = 1$  case. Define a function  $\varphi$  on  $\mathbb{R}$  by setting

$$\varphi(x) = \begin{cases} 1, & 0 \leq x \leq 1, \\ -1, & -1 \leq x < 0, \\ 0, & \text{otherwise.} \end{cases}$$

Then, a direct computation shows that  $\|f\|_{BMO} = 1$ . While the function  $f$  suffices for showing weakly sequential non-completeness of  $BMO$ , we need to smooth it out to treat the  $VMO$  and  $CMO$  cases since  $\varphi \notin VMO$ .

Let  $\rho : \mathbb{R} \rightarrow [0, 1]$  be a smooth even function with compact support, and set  $\rho_\varepsilon(x) = \varepsilon^{-1} \rho(\varepsilon^{-1}x)$ . Fix small  $\varepsilon > 0$  and set  $\varphi_\varepsilon = \rho_\varepsilon * \varphi$ . Then, we have

- $\text{supp } \varphi_\varepsilon \subset [-2, 2]$ ,
- $\|\varphi_\varepsilon\|_{BMO} \sim 1$ ,
- $\varphi_\varepsilon \in VMO$  and hence  $\varphi_\varepsilon \in CMO$  (in view of its compact support).

Now, given  $n \in \mathbb{N}$ , we set  $f_n(x) = \|\varphi_\varepsilon\|_{BMO}^{-1} \cdot \varphi_\varepsilon(x - 10n)$ . Then, we define a subspace  $Y$  of  $BMO$  by

$$Y = \left\{ f = \sum_{n=1}^{\infty} a_n f_n : \mathbf{a} = \{a_n\}_{n \in \mathbb{N}} \in c_0 \right\}.$$

We claim that there exist constants  $C_1, C_2 > 0$  such that

$$C_1 \sup_{n \in \mathbb{N}} |a_n| \leq \left\| \sum_{n=1}^{\infty} a_n f_n \right\|_{BMO} \leq C_2 \sup_{n \in \mathbb{N}} |a_n| \quad (\text{B.3})$$

for any  $\mathbf{a} = \{a_n\}_{n \in \mathbb{N}} \in c_0$ . In fact, it follows from [16, Proposition 3.1.2 (2)], the disjointness of the supports of  $f_n$ , and the fact that the  $L^\infty$ -norm of  $f_n$  is constant in  $n \in \mathbb{N}$  that

$$\left\| \sum_{n=1}^{\infty} a_n f_n \right\|_{BMO} \leq 2 \left\| \sum_{n=1}^{\infty} a_n f_n \right\|_{L^\infty} \leq C_2 \sup_{n \in \mathbb{N}} |a_n|.$$

On the other hand, there exists  $C_1$ , independent of  $k \in \mathbb{N}$ , such that

$$\left\| \sum_{n=1}^{\infty} a_n f_n \right\|_{BMO} \geq \frac{1}{|\text{supp } f_k|} \int_{\text{supp } f_k} |a_k f_k(x)| dx = C_1 |a_k|$$

for any  $k \in \mathbb{N}$ , where we used the fact that  $f_k$  has mean zero in the first step and the second step follows from noting that the size of the support and the  $L^1$ -norm of  $f_k$  do not depend on  $k \in \mathbb{N}$ . This proves (B.3).

From the right-hand side of (B.3), we see that every element in  $Y$  is the limit in  $BMO$  of  $C_c^\infty$ -functions (namely their partial sums). Hence, from [5, Théorème 7 (ii)], we see that  $Y \subset CMO$ . This shows that  $CMO$  contains a copy of  $c_0$ . Hence, from Lemma B.4, we conclude that  $CMO$  is not weakly sequentially complete. By applying Lemma B.2, we also see that  $BMO$  and  $VMO$  are not weakly sequentially complete.  $\square$

We conclude this section by recalling that  $BMO$  is sequentially complete with respect to the weak- $*$  topology as the dual of  $H^1$ . In fact, this is a particular case of the following general result. While the proof is standard, we include it for readers' convenience.

**Lemma B.5.** *Let  $X$  be a Banach space. Then,  $X^*$  is weak- $*$  sequentially complete.*

*Proof.* Let  $\{f_n\}_{n \in \mathbb{N}}$  be a weak- $*$  Cauchy sequence in  $X^*$ . From the Banach-Steinhaus theorem, we see that  $\sup_{n \in \mathbb{N}} \|f_n\|_{X^*} < \infty$ . Then, it follows from the Banach-Alaoglu theorem that there exists a subsequence  $\{f_{n_j}\}_{j \in \mathbb{N}}$  and  $f \in X^*$  such that, as  $j \rightarrow \infty$ ,  $f_{n_j} \rightarrow f$  with respect to the weak- $*$  topology. That is,

$$\langle f, h \rangle = \lim_{k \rightarrow \infty} \langle f_{n_{j_k}}, h \rangle = \lim_{n \rightarrow \infty} \langle f_n, h \rangle$$

for any  $h \in X$ , where the second equality follows from the fact that  $\{f_n\}_{n \in \mathbb{N}}$  is weak- $*$  Cauchy in  $X^*$  (which in particular guarantees uniqueness of the weak- $*$  limit). The proof is complete.  $\square$

**Acknowledgements.** Á.B. acknowledges the support from an AMS-Simons Research Enhancement Grant for PUI Faculty. G.L. and T.O. were supported by the European Research Council (grant no. 864138 “SingStochDispDyn”). The first three authors would like to thank the West University of Timișoara for its hospitality, where part of this paper was prepared.

## REFERENCES

- [1] J. Alvarez, M. Milman,  *$H^p$  continuity properties of Calderón-Zygmund-type operators*, J. Math. Anal. Appl. 118 (1986), no.1, 63–79.
- [2] E. Bednarczuk, K. Leśniewski, *On weakly sequentially complete Banach spaces*, J. Convex Anal. 24 (2017), no. 4, 1341–1356.
- [3] Á. Bényi, G. Li, T. Oh, R.H. Torres, *Compactness of bilinear paraproducts revisited*, preprint.
- [4] Á. Bényi, T. Oh, *Linear and bilinear  $T(b)$  theorems à la Stein*, Proc. Amer. Math. Soc. Ser. B 2 (2015), 1–16.
- [5] G. Bourdaud, *Remarques sur certains sous-espaces de  $BMO(\mathbb{R}^n)$  et de  $bmo(\mathbb{R}^n)$* , Ann. Inst. Fourier (Grenoble) 52 (2002), no. 4, 1187–1218.
- [6] A.P. Calderón, *Intermediate spaces and interpolation, the complex method*, Studia Math. 24 (1964), 113–190.
- [7] M.J. Carro, J. Soria, R.H. Torres, *Extrapolation of compactness for certain pseudodifferential operators*, Rev. Un. Mat. Argentina 66 (2023), no. 1, 177–186.
- [8] R. Coifman, G. Weiss, *Extensions of Hardy spaces and their use in analysis*, Bull. Amer. Math. Soc. 83 (1977), no. 4, 569–645.
- [9] H.O. Cordes, *On compactness of commutators of multiplications and convolutions, and boundedness of pseudodifferential operators*, J. Functional Analysis 18 (1975), 115–131.
- [10] G. David, J.-L. Journé, *A boundedness criterion for generalized Calderón-Zygmund operators*, Ann. of Math. 120 (1984), no. 2, 371–397.

- [11] C. Fefferman, E. Stein,  *$H^p$  spaces of several variables*, Acta Math. 129 (1972), no. 3-4, 137–193.
- [12] H. Feichtinger, *Compactness in translation invariant Banach spaces of distributions and compact multipliers*, J. Math. Anal. Appl. 102 (1984), no. 2, 289–327.
- [13] A. Fragkos, A.W. Green, B.D. Wick, *Multilinear wavelet compact  $T(1)$  theorem*, arXiv:2312.09185 [math.CA].
- [14] M. Frazier, Y.S. Han, B. Jawerth, G. Weiss, *The  $T_1$  theorem for Triebel-Lizorkin spaces*, Harmonic analysis and partial differential equations (El Escorial, 1987), 168–181, Lecture Notes in Math., 1384, Springer, Berlin, 1989.
- [15] M. Frazier, R.H. Torres, G. Weiss, *The boundedness of Calderón-Zygmund operators on the spaces  $\dot{F}_p^{\alpha,q}$* , Rev. Mat. Iberoamericana 4 (1988), no. 1, 41–72.
- [16] L. Grafakos, *Modern Fourier analysis*. Third edition. Graduate Texts in Mathematics, 250. Springer, New York, 2014. xvi+624 pp.
- [17] H. Hanche-Olsen, H. Holden, *The Kolmogorov-Riesz compactness theorem*, Expo. Math. 28 (2010), no. 4, 385–394. Addendum to “The Kolmogorov-Riesz compactness theorem” [Expo. Math. 28 (2010) 385–394]. Expo. Math. 34 (2016), no. 2, 243–245.
- [18] J. Hart, *A new proof of the bilinear  $T(1)$  theorem*, Proc. Amer. Math. Soc. 142 (2014), no. 9, 3169–3181.
- [19] M.A. Krasnosel’skii, *On a theorem of M. Riesz*, Soviet Math. Dokl. 1 (1960), 229–231; translated from Dokl. Akad. Nauk SSSR 131 (1960), 246–248.
- [20] J. Lindenstrauss, L. Tzafriri, *Classical Banach spaces. I. Sequence spaces*. Ergebnisse der Mathematik und ihrer Grenzgebiete [Results in Mathematics and Related Areas], Band 92. Springer-Verlag, Berlin-New York, 1977. xiii+188 pp.
- [21] Y. Meyer, *Les nouveaux opérateurs de Calderón-Zygmund*, (French) [The new Calderón-Zygmund operators] Colloquium in honor of Laurent Schwartz, Vol. 1 (Palaiseau, 1983). Astérisque No. 131 (1985), 237–254.
- [22] Y. Meyer, R. Coifman, *Wavelets. Calderón-Zygmund and multilinear operators*, Translated from the 1990 and 1991 French originals by David Salinger. Cambridge Studies in Advanced Mathematics, 48. Cambridge University Press, Cambridge, 1997. xx+315 pp.
- [23] M. Mitkovski, C.B. Stockdale, *On the  $T_1$  theorem for compactness of Calderón-Zygmund operators*, arXiv:2309.15819 [math.CA].
- [24] J.F. Olsen, P. Villarroya, *Endpoint estimates for compact Calderón-Zygmund operators*, Rev. Mat. Iberoam. 33 (2017), no. 4, 1285–1308.
- [25] K.-M. Perfekt, S. Pott, P. Villarroya, *Endpoint compactness of singular integrals and perturbations of the Cauchy integral*, Kyoto J. Math. 57 (2017), no. 2, 365–393.
- [26] M. Reed, B. Simon, *Methods of modern mathematical physics. I. Functional analysis*. Second edition. Academic Press, Inc. [Harcourt Brace Jovanovich, Publishers], New York, 1980. xv+400 pp.
- [27] E. Stein, *Singular integrals and differentiability properties of functions*, Princeton Mathematical Series, No. 30. Princeton University Press, Princeton, NJ, 1970. xiv+290 pp.
- [28] E. Stein, *Harmonic analysis: real-variable methods, orthogonality, and oscillatory integrals*, Princeton Mathematical Series, 43. Monographs in Harmonic Analysis, III. Princeton University Press, Princeton, NJ, 1993. xiv+695 pp.
- [29] A. Uchiyama, *On the compactness of operators of Hankel type*, Tohoku Math. J. 30 (1978), no. 1, 163–171.
- [30] P. Villarroya, *A characterization of compactness for singular integrals*, J. Math. Pures Appl. 104 (2015), no. 3, 485–532.
- [31] P. Villarroya, *A global  $T_b$  theorem for compactness and boundedness of Calderón-Zygmund operators*, J. Math. Anal. Appl. 480 (2019), no. 1, 123323, 41 pp.
- [32] K. Yosida, *Functional analysis*, Reprint of the sixth (1980) edition. Classics in Mathematics. Springer-Verlag, Berlin, 1995. xii+501 pp.

ÁRPÁD BÉNYI, DEPARTMENT OF MATHEMATICS, 516 HIGH ST, WESTERN WASHINGTON UNIVERSITY, BELLINGHAM, WA 98225, USA.

*Email address:* benyia@wwu.edu

GUOPENG LI, SCHOOL OF MATHEMATICS, THE UNIVERSITY OF EDINBURGH, AND THE MAXWELL INSTITUTE FOR THE MATHEMATICAL SCIENCES, JAMES CLERK MAXWELL BUILDING, THE KING’S BUILDINGS, PETER GUTHRIE TAIT ROAD, EDINBURGH, EH9 3FD, UNITED KINGDOM

*Email address:* guopeng.li@ed.ac.uk

TADAHIRO OH, SCHOOL OF MATHEMATICS, THE UNIVERSITY OF EDINBURGH, AND THE MAXWELL INSTITUTE FOR THE MATHEMATICAL SCIENCES, JAMES CLERK MAXWELL BUILDING, THE KING'S BUILDINGS, PETER GUTHRIE TAIT ROAD, EDINBURGH, EH9 3FD, UNITED KINGDOM

*Email address:* `hiro.oh@ed.ac.uk`

RODOLFO H. TORRES, DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CALIFORNIA, 900 UNIVERSITY AVE., RIVERSIDE, CA 92521, USA

*Email address:* `rodolfo.h.torres@ucr.edu`